

Understanding Crohn's disease and ulcerative colitis



Table of contents

Introduction to inflammatory bowel disease (IBD)	1
Who develops IBD?	1
Causes	2
Risk factors	2
Diagnosis	2
What you need to know about Crohn's disease and ulcerative colitis	3
Crohn's disease	3
Ulcerative colitis	3
Living With IBD	
Staying on track with treatment	4
Lifestyle changes	5
Physical activity	6
Emotional health	6
Proper nutrition	7
Surgery	9
Pregnancy	10
Ongoing care	10
References	12
Pasourcas	1/

This publication is for informational purposes only. It is not intended to be a substitute for professional medical advice, diagnosis or treatment. Always seek the advice of your physician or other qualified healthcare provider with any questions you may have regarding a medical condition. Never disregard professional medical advice or delay in seeking it because of something you have read in this publication. If you think you may have a medical emergency, call your physician or 911 immediately. Walgreens Specialty Pharmacy does not recommend or endorse any specific tests, physicians, products, procedures, opinions or other information that may be mentioned in this publication. Reliance on any information provided in this publication is solely at your own risk.

This publication was created by and is provided as a service of Walgreens Specialty Pharmacy.

Introduction to inflammatory bowel disease

Learning how to manage your inflammatory bowel disease (IBD) might feel like a challenge, especially at first. But understanding your diagnosis can help you take control of your health. There is no cure for IBD. But a number of treatment options can help you manage your symptoms. This booklet provides information about IBD, what to expect after diagnosis and how to manage symptoms to live a full and active life.


IBD refers to chronic, or long-term, conditions that affect the gastrointestinal (GI) tract. These conditions can get better and worse over time. The GI tract is the part of the body that digests food. It also absorbs nutrients and expels waste. The GI tract includes the mouth, esophagus, stomach, small intestine, large intestine and anus.

The two types of IBD are Crohn's disease and ulcerative colitis. Both cause inflammation in the GI tract. Crohn's disease can affect any part of the GI tract and all layers of the intestines. Ulcerative colitis affects just the lining of the large intestine, which includes the colon and rectum.^{1,2}

Who develops IBD?

In the United States, an estimated 3.1 million adults are diagnosed with IBD.³ It can occur at any age, but people are usually diagnosed between the ages of 15 and 35. Although males and females can both have IBD, females tend to have a predisposition for developing IBD based on the X chromosome.^{4,5}

The GI tract



Causes

The exact cause of IBD is not known. Experts believe it is caused by a number of factors, including⁴:

- Environment
- Genes, which decide inherited traits such as eye color
- Immune system problems

Normally, the immune system seeks out and attacks intruders, like bad bacteria or viruses that can cause illness. It also helps the body recover from sickness. But it does not always work properly. In people with IBD, the immune system mistakenly attacks harmless bacteria, food and other materials in the intestine. This can cause swelling in the GI tract.

Risk factors

Risk factors for IBD include4:

- Antibiotics, which can affect levels of bacteria in the GI tract
- Appendectomy, or surgery to remove the appendix (risk factor for Crohn's disease)
- Cigarette smoking (risk factor for Crohn's disease)
- Family history
- Nonsteroidal anti-inflammatory drugs (NSAIDs) such as aspirin, ibuprofen or naproxen

Some studies suggest that diet, use of hormones and certain medications, stress and low levels of vitamin D might raise the risk of IBD.^{6,7}

Diagnosis

Your healthcare provider likely diagnosed your IBD based on your symptoms, a physical exam and tests.^{8,9} Diagnostic tests for IBD often include:

- Blood tests
- X-rays or other body scans
- Analysis of stool samples for bleeding or infection

Your healthcare provider probably also viewed your GI tract using procedures such as^{8,9}:

- Biopsy. A small piece of tissue is removed from the affected part of the GI tract for testing.
- Colonoscopy. A thin, flexible tube with a camera is used to view the entire colon.
- Flexible sigmoidoscopy. A thin, flexible tube with a camera is used to view the end of the colon.
- Upper endoscopy. A thin, flexible tube with a camera is used to view the upper portion of the digestive system, which includes the stomach and small intestine.





Understanding your diagnosis can help you better manage your health. Crohn's disease and ulcerative colitis are long-term conditions with no cure. But, starting treatment soon after diagnosis can help manage your symptoms so you can live a full and active life.⁴

Crohn's disease

Who develops Crohn's disease?

Crohn's disease affects men and women equally. It is more often diagnosed among people ages 20 to 30.4,10

Signs and symptoms

Crohn's disease is a long-term condition. It can get better and worse over time. While there is no cure, medication can control symptoms. Crohn's disease is active when symptoms are present. This is also called a flare. Remission is when symptoms stop. Symptoms of Crohn's disease can differ for each patient. Symptoms can range from mild to severe.

Common GI symptoms include1:

- Abdominal pain and cramps
- Constipation
- Diarrhea
- Feeling of incomplete bowel movements
- Rectal bleeding
- Urgent need to have a bowel movement

Other symptoms of Crohn's disease might also include¹:

- Fatigue (tiredness)
- Fever
- Menstrual changes
- Night sweats
- Poor appetite
- Weight loss

Ulcerative colitis

Who develops ulcerative colitis?

Ulcerative colitis can occur at any age, though most people are diagnosed in their mid-30s.¹¹ It is more common in men than in women. Men are also more likely than women to be diagnosed with ulcerative colitis in their 50s and 60s.⁴

Signs and symptoms

Like Crohn's disease, ulcerative colitis is a long-term condition. It can get better and worse over time. While there is no cure, medication can control symptoms.

Symptoms of ulcerative colitis can include²:

- Abdominal pain and cramps
- Delayed growth in children
- Diarrhea
- Fatigue
- Loose, bloody stools
- Poor appetite
- Urgent need to have a bowel movement
- Weight loss

28 Living with IBD

There is no cure for IBD. But there are many ways to treat the symptoms. A healthcare provider who specializes in GI problems, known as a gastroenterologist, can help direct your treatment.

Treatment can help in achieving a range of goals, including:

- Decreasing inflammation in the GI tract
- Improving quality of life
- Inducing remission or stopping symptoms
- Maintaining remission or preventing flares
- Reducing risk for complications in the GI tract or other parts of the body

Work with your healthcare provider to find the best treatment for you. Your healthcare provider might ask you to keep track of symptoms in a diary. For example, you might log how many and what type of bowel movements you have each day. Or, you might track your weight.

Treatment can control symptoms. But there will be times when symptoms return. These flares can be triggered by many things, including¹²:

- Certain foods
- Certain medications, like NSAIDs or antibiotics
- Cigarette smoking (for patients with Crohn's disease)
- Not taking medication as prescribed
- Stress

IBD can also cause other complications. These can occur in the GI tract or other parts of the body. Complications might include⁴:

- · Abscess, or collection of pus in the GI tract
- Anemia that can cause fatigue

- · Arthritis, or swollen joints
- Bone problems, like weak bones
- Colon cancer
- Eye problems, like swelling or irritation
- Fistulas, or abnormal tunnels in the intestines caused by ulcers
- Kidney stones
- Liver disease
- Malabsorption and malnutrition that could cause vitamin or mineral deficiencies or affect growth in children
- Mouth sores
- Narrowed intestine that can cause blockage
- Perforated bowel, or holes in the intestinal lining
- Skin problems, like sores, ulcers or rashes
- Toxic megacolon, a dangerous and rapid widening of the colon
- Ulcers in the GI tract

Some complications might require urgent care. Be prepared. Ask your healthcare provider about IBD complications before they occur. Find out when you should seek immediate medical care.

Staying on track with treatment

Medication can help manage IBD symptoms. The main types of drugs used to treat IBD are⁴:

- Aminosalicylates
- Antibiotics
- Biologic therapies
- Corticosteroids
- Immunomodulators



There is not one single treatment that works for all people with IBD. Treatments are prescribed specifically for each individual. Work with your healthcare provider to find a treatment that works best for you. It might change over time. Your healthcare provider can adjust your prescribed treatment as needed.

No matter what medications are chosen for your treatment, you should take them exactly as prescribed by your healthcare provider — at the right times and at the correct doses. Do not stop taking your medications without asking your healthcare provider first. If your treatment routine starts to feel too hard, ask your healthcare provider or pharmacist for help managing your medications. Staying on track with treatment is key to managing symptoms and improving your health.

Lifestyle changes

Along with medication, lifestyle changes can help you manage IBD.^{13,14} Some lifestyle changes might be small changes to your daily routine. For example, you might start to locate restrooms before an outing. Or, you might start bringing extra underwear and toilet paper wherever you go. There are also apps, such as "Flush," which uses your GPS to find the nearest public restroom.¹⁵

Other lifestyle changes might be larger goals. For example, smoking can make Crohn's disease worse.⁴ It can also cause a host of other health problems. Quitting smoking can lessen GI symptoms and improve your health. If you smoke, talk to your healthcare provider about ways to stop.



Habits that help you stay active, manage stress and eat right can help you feel better each day. These changes can also help improve IBD symptoms.

Physical activity

Staying active is good for your health. It can help you keep a healthy weight. This can reduce strain on your joints. It might also help with joint problems caused by IBD. Exercise is also a good way to relieve any stress that could make IBD symptoms worse.¹³ Many types of activity might be useful, including:

- Aerobic activity, like walking or swimming
- Strengthening exercises, like weight training
- Stretching and range-of-motion routines, like yoga or tai chi

Talk to your healthcare provider about what types of exercise might be right for you.

Emotional health

Sometimes, living with IBD can feel difficult. You might feel angry, anxious or embarrassed. This is normal. But it might make things feel more stressful than usual. Stress can make it harder to deal with the demands of living with IBD. It might also trigger symptoms. Take steps to control your stress:

- Find out what causes your stress. Try keeping a journal or diary to pinpoint the sources.
- Try to avoid things that cause you stress.
- Find positive ways to cope. Make time for things you enjoy. Relax in a quiet space each day.

You might also feel sad or depressed about your condition. This is normal, too. These feelings might be worse when you are first diagnosed, in the early stages of the disease or during a flare.

It's important to know the symptoms of depression and when to get help. Symptoms of depression can include¹⁶:

- Aches and pains that don't go away with treatment
- Being tired or lacking energy
- · Eating too much or too little
- Feeling hopeless or negative
- Feeling restless or irritable
- Feeling sad, empty or anxious most of the time
- · Feeling worthless, helpless or guilty
- Having a hard time concentrating or making decisions
- Losing interest or pleasure in things you used to enjoy
- Sleeping too much or too little
- · Thinking about death or suicide

If you have thoughts of suicide, call 911 or your local emergency services number. You can also call a healthcare provider, mental health professional, crisis center or hotline for help. You can also can text or call the suicide hotline number, which is 988. The crisis line text is 741741.^{17,18}

Children and teens might express depression in additional ways, including¹⁹:

- Extreme sensitivity to failure
- Frequent complaints of headaches or stomachaches
- Increased anger
- Low energy or constant boredom
- Low self-esteem
- Poor performance or attendance at school
- Social isolation
- Talk of running away from home

Talk with your healthcare provider or your child's healthcare provider about depression. Your healthcare provider might suggest counseling, an antidepressant medication or both. You might also want to seek support from friends and family. Or, consider joining a support group or online message board where you can talk to others with IBD.

Proper nutrition

Poor nutrition is common in patients with IBD. This is especially true in Crohn's disease. This is because the intestines are less able to digest and absorb nutrients from food. IBD symptoms can also make it hard to eat well all the time. But proper nutrition is key for good health and normal growth. This is especially vital for children with IBD.²⁰ Talk with your healthcare provider about how to improve your nutrition.



Your healthcare provider might refer you to a dietitian. Together, you can build a diet that works for you. In general, it is most important to eat a well-balanced diet and drink enough fluids to stay hydrated. There is no single list of foods to eat or avoid. Certain foods might bother you but not other people with IBD. This can change over time.

You might also need to eat differently at certain times. For example, it might be best to eat smaller meals more frequently during flares. Five small meals (fist-sized portions) every three or four hours is easier on your digestive system than three large meals.¹⁴

Try keeping a food diary to track what you eat. You can also log symptoms and when they occur. Your notes might help you see which foods or meal times cause symptoms. Your log can also help your healthcare provider or dietitian see if you are getting enough nutrients from your diet.

Some people with Crohn's disease have a narrowed part of the small intestine. For these people, a low-fiber with low-residue diet might help reduce symptoms. This diet avoids certain foods — like raw produce, nuts and seeds — that leave undigested material to pass through the intestines.



This can reduce the number and size of stools. It is typically only used for a short time, until symptoms improve.¹⁴ Ask your healthcare provider if this diet might help you.

Bone health

Steroids are one type of medication used to treat IBD. If used for a long time, they can cause osteoporosis, or weak bones.⁴ This is true in women and men. Keep track of your bone health if you are taking steroids. Your healthcare provider might want you to take a bone density test. This test looks at the strength of your bones. It can also assess your risk of breaking a bone. You can take other steps to help prevent weak bones²¹:

 Get enough calcium and vitamin D in your diet. Talk to your healthcare provider about the right amount and how to reach that level.

- Do not smoke cigarettes.
- Engage in regular physical activity, as directed by your healthcare provider.
- · Limit or avoid alcohol.

In children, long-term steroid treatment can also delay growth. Children are sometimes treated with steroids every other day to reduce this effect.²² Ask your child's healthcare provider about protecting your child's bones.

Surgery

Even with medication and lifestyle changes, some people will need surgery to control IBD symptoms. About 70% of people with Crohn's disease will need surgery. Among those with ulcerative colitis, up to one-third will need surgery.⁴ Table 3 lists common surgical procedures for IBD

Table 3. Surgical procedures for IBD^{23,24}

Procedure	Description
Crohn's disease	
Colectomy/proctocolectomy with ileostomy	 Colectomy removes all or part of the colon Proctocolectomy removes the colon and rectum Ileostomy creates an opening through the abdomen that allows waste to empty out of the body into an attached pouch
Fistula or abscess repair	 Fixes abnormal connections between different parts of the GI tract Removes abscesses
Resection	Removes affected portions of the intestine and reconnects healthy sections of intestine
Strictureplasty	Widens narrowed parts of intestine
Ulcerative colitis	
Proctocolectomy with ileostomy	Removes the colon and rectum and creates an opening through the abdomen that allows waste to empty out of the body into an attached pouch
Restorative proctocolectomy or ileal pouch anal anastomosis	Removes the colon and rectum and creates a path for waste to empty through the anus



Pregnancy

Active IBD and past pelvic surgery to treat IBD might affect a woman's chances of getting pregnant. The risk of complications rises when IBD is active during conception and pregnancy. Ideally, women with IBD should be in remission for at least six months before getting pregnant.⁴

Some medications for IBD can affect fertility in both men and women. Certain IBD medications are not safe for pregnancy. Tell your healthcare provider if you are planning to get pregnant or if you get pregnant while taking IBD medications. Your healthcare provider might need to change your medication. Women with IBD who plan to breastfeed should also ask a healthcare provider or pharmacist about which medications are safe to take while nursing.²⁵

Ongoing care

Work with your healthcare provider to keep track of your progress and your symptoms. You can discuss how well your medication and lifestyle changes are working. Your healthcare provider can also monitor side effects and adjust your treatment as needed. Each visit can help you stay on track with treatment and better manage your IBD.

We provide this information because the more you know about IBD, the better you'll be able to manage it.

Additionally, the Walgreens Specialty360
Therapy Team is here to support you with
dependable, personalized service to help manage
your medication side effects and stay on track
with your prescribed therapy.


We look forward to being a member of your healthcare team and helping you get the best results from your treatment.



References

- What is Crohn's disease? Crohn's and Colitis Foundation of America Web site. https://www.crohnscolitisfoundation.org/ patientsandcaregivers/what-is-crohns-disease. Accessed February 21, 2024.
- 2. What is ulcerative colitis? Crohn's and Colitis Foundation of America Web site. https://www.crohnscolitisfoundation.org/patientsandcaregivers/what-is-ulcerative-colitis. Accessed February 21, 2024.
- 3. Centers for Disease Control and Prevention. Prevalence of IBD. CDC Web site. https://www.cdc.gov/ibd/data-and-statistics/prevalence.html. Accessed February 21, 2024.
- 4. The facts about inflammatory bowel diseases. Crohn's and Colitis Foundation of America Web site. https://www.crohnscolitisfoundation.org/sites/default/files/legacy/assets/pdfs/ibdfactbook.pdf. Published November 2014. Accessed February 21, 2024.

- National Library of Medicine. Impact of Female Gender in Inflammatory Bowel Diseases: A Narrative Review. National Library of Medicine Web site. https://www.ncbi.nlm.nih.gov/pmc/ articles/PMC9958616/#:~:text=Females%20 in%20familial%20IBD%20population,female%20 to%20female%20transmission%20vs. Accessed February 24, 2024.
- Ananthakrishnan AN. Environmental risk factors for inflammatory bowel disease. Gastroenterol Hepatol (NY). 2013;9(6):367-374. http://www. ncbi.nlm.nih.gov/pmc/articles/PMC3736793/. Accessed February 21, 2024.
- 7. Crockett SD, Porter CQ, Martin CF, Sandler RS, Kappelman MD. Isotretinoin use and the risk of inflammatory bowel disease: A case control study. Am J Gastroenterol. 2010;105(9):1986-1993. http://www.nature.com/ajg/journal/v105/n9/full/ajg2010124a.html. Accessed February 21, 2024.



- Crohn's diagnosis and testing. Crohn's and Colitis
 Foundation of America Web site. https://www.
 crohnscolitisfoundation.org/what-is-crohnsdisease/diagnosis-testing. Accessed
 February 21, 2024.
- Ulcerative colitis. National Institute of Diabetes and Digestive and Kidney Diseases Web site. https://www.niddk.nih.gov/health-information/ digestive-diseases/ulcerative-colitis/diagnosis. Updated September 2020. Accessed February 21, 2024.
- Crohn's & Colitis Foundation. Overview of Crohn's Disease. Crohn's & Colitis Foundation Web site. https://www.crohnscolitisfoundation.org/ patientsandcaregivers/what-is-crohns-disease/ overview. Accessed February 24, 2024.
- Crohn's & Colitis Foundation. Overview of Ulcerative Colitis. Crohn's & Colitis Foundation Web site. https://www.crohnscolitisfoundation. org/patientsandcaregivers/what-is-ulcerativecolitis/overview. Accessed February 24, 2024.
- Managing flares and other IBD symptoms. Crohn's and Colitis Foundation of America Web site. http://www.crohnscolitisfoundation.org/ assets/pdfs/Managing-flares.pdf. Updated June 2019. Accessed February 21, 2024.
- Neeraj N, Fedorak RN. Exercise and inflammatory bowel disease. Can J Gastroenterol. 2008;22(5):497-504. http://www.ncbi.nlm.nih. gov/pmc/articles/PMC2660805/. Accessed February 21, 2024.
- 14. Inflammatory bowel disease and irritable bowel syndrome: similarities and differences. Crohn's and Colitis Foundation of America Web site. http://www.crohnscolitisfoundation.org/assets/ pdfs/ibd-and-irritable-bowel.pdf. Published July 2014. Accessed February 21, 2024.
- Apple. Flush Toilet Finder & Map. Apple App Store Web site. https://apps.apple.com/us/app/ flush-toilet-finder-map/id955254528. Accessed February 24, 2024.
- 16. Depression: signs and symptoms. National Institute of Mental Health Web site. https:// www.nimh.nih.gov/health/topics/depression/ index.shtml#part_145397. Updated April 2023. Accessed February 21, 2024.

- Suicide & Crisis Lifeline. 988 Suicide & Crisis Lifeline. Suicide & Crisis Lifeline Web site. https://988lifeline.org/. Accessed February 24, 2024.
- 18. Crisis Text Line. Need to vent? Crisis Text Line Web site. https://www.crisistextline.org/. Accessed February 24, 2024.
- Depression in children and teens. American Academy of Child and Adolescent Psychiatry Web site. https://www.aacap.org/AACAP/ Families_and_Youth/Facts_for_Families/FFF-Guide/The-Depressed-Child-004.aspx. Updated October 2018. Accessed February 21, 2024.
- Lucendo AJ, Rezende LC. Importance of nutrition in inflammatory bowel disease.
 World J Gastroenterol. 2009;15(7):2081-2088. http://www.ncbi.nlm.nih.gov/pmc/articles/ PMC2678578/. Accessed February 21, 2024.
- 21. Bone loss in IBD. Crohn's and Colitis Foundation of America Web site. https://www.crohnscolitisfoundation.org/sites/default/files/2020-03/boneloss.pdf. Published May 1, 2012. Accessed February 21, 2024.
- 22. Deshmukh CT. Minimizing side effects of systemic corticosteroids in children. Indian J Dermatol Venereol Leprol. 2007;73:218-221. http://www.ijdvl.com/text.asp?2007/73/4/218/33633. Accessed February 21, 2024.
- 23. Inflammatory bowel disease fact sheet. Office on Women's Health Web site. https://www.womenshealth.gov/publications/our-publications/fact-sheet/inflammatory-bowel-disease.html. Updated June 7, 2022. Accessed February 21, 2024.
- 24. Surgery for ulcerative colitis. Crohn's and Colitis Foundation of America Web site. https://www.crohnscolitisfoundation.org/what-is-ulcerative-colitis/surgery. Accessed February 21, 2024.
- 25. Inflammatory bowel disease. Organization of Teratology Information Specialists Mother to Baby Web site. http://mothertobaby.org/fact-sheets/inflammatory-bowel-disease-pregnancy/pdf/. Updated January 1, 2022. Accessed February 21, 2024.

Resources

You might find it helpful to contact these organizations for additional support and resources.*

American College of Gastroenterology (ACG)

www.gi.org/topics/inflammatory-bowel-disease 301-263-9000 www.facebook.com/AmCollegeGastro/ @AmCollegeGastro

The ACG is a clinical organization of digestive health specialists. The ACG website features information on GI conditions, including IBD, in both English and Spanish. The site also provides resources for finding a gastroenterologist and other patient support.

American Society of Colon and Rectal Surgeons (ASCRS)

www.fascrs.org/patients 847-290-9184 www.facebook.com/fascrs @fascrs_updates

The ASCRS is a group of doctors who specialize in treating patients with conditions that affect the colon, rectum and anus. The group's website provides disease information, a patient resource library and a directory of surgeons.

Crohn's & Colitis Foundation of America (CCFA)

www.ccfa.org 800-932-2423 www.facebook.com/ccfafb @CrohnsColitisFn

The CCFA is a nonprofit, volunteer organization devoted to finding a cure for Crohn's disease and ulcerative colitis. The CCFA website includes information on treatment, nutrition, surgery options and doctors. Brochures, webcasts, information on clinical trials and links to other organizations are also available. The companion website, www.ibdetermined.org, provides links to more information and ways to connect with others through social media and a smartphone app.

MedlinePlus

www.nlm.nih.gov/medlineplus/crohnsdisease.html (Crohn's disease)

www.nlm.nih.gov/medlineplus/ulcerativecolitis.html (ulcerative colitis)

www.facebook.com/mplus.gov/ @medlineplus

MedlinePlus is a comprehensive online health resource from the U.S. National Library of Medicine. Its website features educational materials on diagnosis, treatment and support for IBD. It also features links to the latest news and clinical trials on IBD and other GI conditions.

^{*}The referenced organizations are provided for informational purposes only. They are not affiliated with, and have not provided funding to, Walgreens Specialty Pharmacy for this booklet. Walgreens Specialty Pharmacy does not endorse or recommend any specific organization.



Notes		



Notes		

